

## Non-Immune Rabbit Immunoglobulin

60-0060 6 mL Non-Immune Rabbit Immunoglobulin, Ready-To-Use  
Concentration: See container label

### Intended Use

For In Vitro Diagnostic Use.

This product is intended to be used in normal and neoplastic formalin fixed paraffin embedded (FFPE) tissue sections in immunohistochemical (IHC) detection methodology. Interpretation must be made within the context of the patient's clinical history and other diagnostic test by a qualified pathologist.

### Description

This non-immune rabbit immunoglobulin is derived from non-immune rabbits and has no known specificity. The concentration of rabbit immunoglobulin in this reagent is similar to that of many primary rabbit antibodies used for immunohistostaining. No background or specific staining was observed when tested on human tissues using Genemed Acu-Stain™ HRP Detection Kits. This product may be used as a negative control for qualitative assessment of background staining for primary rabbit antibodies in IHC staining of sections of routine FFPE human tissue.

### Reagent provided

This product contains 1 µg/mL of normal (non-immune) rabbit immunoglobulin diluted in 10 mM phosphate buffered saline (PBS), pH 7.2 containing 1% bovine serum albumin (BSA) and 0.09% sodium azide (NaN<sub>3</sub>) as antimicrobial agent.

### Precautions

For professional users.

Proper handling of this product as with any product derived from biological sources according to local and applicable regulations.

Sodium azide is a toxic chemical. The concentration in this product is not classified as hazardous, however, the build-ups of NaN<sub>3</sub> may react with lead and copper plumbing to form highly explosive metal azides. Flush the disposed reagent with large volume of water to prevent azide build-up.

### Usage

#### Dilution

60-0060: Ready-To-Use

#### Staining procedure

Incubate this non-immune rabbit antibody with tissue section for 30-60 minutes at room temperature. Follow the instructions from the selected detection system.

#### Epitope retrieval:

Use the same condition as rabbit primary antibody under testing.

#### Staining pattern

No staining

### Storage

Store at 2-8°C.

### References

1. Elias JM, et al. Am J Clin Pathol. 1989 Dec;92(6):836-43.
2. Diagnostic Immunohistochemistry by David J Dabbis, 3<sup>rd</sup> Ed. 2010

### Symbols



Catalog No.



Batch No.



In Vitro Diagnostic Use



Temperature Range



Use By

